

## IN THE CLAIMS:

Amend Claim 18 to read:

Claim 18 (amended): An assay for determining the concentration of epidermal growth factor receptor in a biological sample from a female patient, the assay comprising:

- a) obtaining a biological sample from the female;
- b) contacting an amount of a first purified antibody that specifically reacts with a first epitope of the extracellular ligand binding domain of sErbB1 with the biological sample to be tested, wherein the first purified antibody is modified with a first labeling moiety;
- c) contacting the sample with an amount of a second purified antibody that specifically reacts with a second epitope of the extracellular ligand binding domain of sErbB1, wherein the second purified antibody is modified with a second labeling moiety, and wherein the second purified antibody does not competitively inhibit the binding of the first purified antibody;
- d) detecting the co-presence of the first and second labels to determine the concentration of the epidermal growth factor receptor complexed with the antibodies;  
  
wherein one of the antibodies is chosen from the group consisting of: MAb R.1 and antibodies which competitively inhibit the binding of MAb R.1 to ErbB1; and wherein the other antibody is chosen from the group consisting of MAb 528 and antibodies which competitively inhibit the binding of MAb 528 to ErbB1
- e) comparing the concentration of soluble epidermal growth factor receptor obtained in step d) with a normal value; and

f) correlating a decrease in the concentration of soluble epidermal growth factor receptor in the biological sample with the presence of an ovarian carcinoma in the patient.